

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

What is the effect of a charge between light and heavy cavalry, the light cavalry having the greater energy and the heavy the greater momentum?

Proposed by DE VOLSON WOOD, M. A., M. So., C. E., Professor of Engineering, Stevens Institute
of Technology, nioboken, New Jersey.

A hollow sphere filled with frictionless water rolls down a rough plane whose length is l and inclination θ ; when half way down the water suddenly freezes and adheres to the sphere. Required the time of the descent.

 Proposed by ALFRED HUME, C. E., D. So.. Professor of Mathematics in the University of Mississippi.

A heavy bar AB of length a falls about its lower end B from a vertical to a horizontal position, when the end A is suddenly fixed and B is set free, so that the bar falls into a vertical position AB as at first; then A is set free, and B is fixed, so that the bar again falls about B into a horizontal position, when the end A is suddenly fixed, and B is set free, and so on; find the angular velocity ω of the bar about the upper end, when it takes a vertical position for the nth time.

[Selected from Price's Infinitesimal Calculus.]

SOLUTION TO THE CELEBRATED INDETERMINATE EQUATION.

$$x^2 - Ny^2 = \pm 1.$$

To obtain the values of x and y, in integers, without taking the square root of N, by continued fractions. On account of requiring a certain class of non-quadrate numbers for N, to satisfy the -1, and causing more or less confusion, this part is set aside for future time. The +1 is found in all values of N, and our equation becomes $x^2 = Ny^2 + 1$.

Then
$$Ny^2+1=\Box=\left\lfloor\frac{mNy}{n}-1\right\rfloor^2$$
 when reduced easily gives $y=\frac{2mn}{m^2}\frac{2mn}{N-n^2}$, and $x=\frac{mNy}{n}-1,\ldots$ (A)

Let $\frac{m}{n}=\frac{1}{2}$, then $y=\frac{4}{N-4}$. Let $\frac{N=2,3,\Box,5,6,8,\Box}{x=3\ 7\ 9\ 5\ 3}$

$$\frac{m}{n}=\frac{1}{3}\text{ , and }y=\frac{6}{N-9},\frac{N=3,7,8,10,11,12,(13),}{y=1\ 3\ 6\ 6\ 3\ 2}$$

$$x=2\ 8\ 17\ 19\ 10\ 7$$

$$\frac{m}{n}=\frac{1}{4}\text{ , and }y=\frac{8}{N-16},\frac{N=14,15,\Box,17,18,(19),20,}{y=4\ 8\ 8\ 4\ 2,}$$

$$x=15\ 31\ 33\ 17\ 9$$

$$\frac{m}{n}=\frac{1}{5}\text{ , }y=\frac{10}{N-25},\frac{N=15,(21),(22),23,24,\Box,26,27,(28),(29),30,}{y=1\ 5\ 10\ 10\ 5\ 2}$$

$$x=4\ 24\ 49\ 51\ 26\ 11$$

$$\frac{m}{n}=\frac{1}{6}\text{ , }y=\frac{12}{N-36},\frac{N=24,(31),32,33,34,35,\Box,37,38,39,40,(41),42,}{y=1\ 3\ 4\ 6\ 12\ 12\ 6\ 4\ 3\ 2}$$

$$x=5\ 17\ 23\ 35\ 71\ 73\ 37\ 25\ 19\ 13$$

$$\frac{m}{n} = \frac{1}{7} , \qquad y = \frac{14}{N-49}, \frac{N=35,(43),(44),(45),(46),47,48, \square, 50,51,(52),(53),}{y=1}, \frac{7}{14}, \frac{14}{14}, \frac{7}{48}, \frac{7}{99}, \frac{50,51}{99}, \frac{50,51,51}{99}, \frac{50,51}{99}, \frac{50,51,51}{99}, \frac{50,51,51,51}{99}, \frac{50,51$$

It is seen that the position of the numbers N, between the two consecutive square numbers, determines the ease of obtaining the values of x and y. As, let a^2 and a_2^2 , represent the two squares, then in $a^2+a,y=2$, in numbers N, of the form of $a^2+2a,y=1$. The more complex numbers are always found midway between a^2+a and a^2+2a .

There are two values of x and y, N being of the form of a^z-1 . It is, however, the second higher value of x and y, as x_z and y_z .

Other series can easily be found giving, the values also in a regular series.

A. H. BELL, Hillsboro, Illinois.

(TO BE CONTINUED.)

AVERAGE AND PROBABILITY.

Conflucted by B. F. FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.

Solution to a Problem by Professor G. B. M. ZERR, A. M., Principal of Schools, Staunton, Virginia.

Three persons A., B., C., throw with three dice. They each stake \$10, and the one who first throws exactly 10 with the three dice, takes the whole stake. Find the expectation of each.

There is probably no subject that is more interesting than the subject of Average and Probability, and that part of this subject that deals